

ETHIODEMOGRAPHY AND HEALTH

CHAPTER 3

POPULATION AND ENVIRONMENT

Ethiopia Focus with a Global Lens

Aynalem Adugna, PhD

EthioDemographyAndHealth.org

Revised October 2026

APA 7TH EDITION | CITATION EXAMPLES

Reference-list entry

Adugna, A. (2026). Population and environment: Ethiopia focus with a global lens [Unpublished chapter manuscript]. EthioDemographyAndHealth.org.

CHAPTER 3 | TABLE OF CONTENTS

List of Acronyms and Abbreviations	3
PART I — From Stockholm to Addis Ababa, 1972–2027	4
3.1 Concepts, pathways, and metrics.....	5
3.2 Demographic change and environmental pressures	6
3.3 Land-use and land-cover change, forests, and restoration	7
3.4 Agriculture, pastoral systems, and rangeland dynamics	8
Part I synthesis and evidence audit.....	8
PART II — Water, Climate, Food Security, and Environmental Health	9
3.5 Water resources, irrigation, and watersheds.....	9
3.6 Climate variability and change: exposure, sensitivity, and adaptation	10
3.7 Food security, nutrition, and environmental shocks.....	10
3.8 Urbanization, air quality, and environmental health	11
Part II synthesis: An Ethiopian environmental-health agenda for COP32.....	12
PART III — Energy, Biodiversity, Disaster Risk, and Infrastructure.....	12
3.9 Energy transitions and household fuels	12
3.10 Biodiversity, protected areas, and human pressure	13
3.11 Disasters, hazards, and ecosystem services	14
3.12 Infrastructure, mining, and environmental footprints.....	14
Part III synthesis: Four connected transitions	15
PART IV — Environmental Governance and the Road to COP32.....	16
3.13 Waste, sanitation, and circularity	16
3.14 Gender, equity, and environmental justice	16
3.15 Governance, policy, and financing for sustainability	17
3.16 Data, methods, and indicators	18
3.17 Case studies and Ethiopian successes.....	19
Part IV synthesis: From hosting to implementation	20
Chapter 3 references	20

CHAPTER 3 | List of Acronyms and Abbreviations

Acronym	Description
CHIRPS	Climate Hazards Center InfraRed Precipitation with Station data
CO₂	Carbon dioxide
COP	Conference of the Parties (to the UNFCCC)
COP32	Thirty-second Conference of the Parties to the UNFCCC (Addis Ababa, 2027)
EIA	Environmental impact assessment
ERA5	Fifth-generation atmospheric reanalysis produced by ECMWF
FAO	Food and Agriculture Organization of the United Nations
GHG	Greenhouse gas
GIS	Geographic information system
IPAT	Impact = Population × Affluence × Technology (conceptual identity)
IPC	Integrated Food Security Phase Classification
IPCC	Intergovernmental Panel on Climate Change
LULC	Land use and land cover
MRV	Measurement, reporting, and verification
NAP	National Adaptation Plan
NDC	Nationally Determined Contribution
NDVI	Normalized Difference Vegetation Index
PM_{2.5}	Fine particulate matter with aerodynamic diameter ≤ 2.5 micrometres
SDG	Sustainable Development Goal
UNEP	United Nations Environment Programme
UNFCCC	United Nations Framework Convention on Climate Change
WASH	Water, sanitation, and hygiene
WHO	World Health Organization

PART I. Opening perspective | From Stockholm to Addis Ababa, 1972–2027

The global environmental agenda has expanded since the first global environmental conference in 1972, rather than simply changing subjects. The Stockholm Conference of 1972 made pollution, human health, and the protection of the natural environment matters of international cooperation. Rio de Janeiro in 1992 connected these concerns to development, biodiversity, forests, and the emerging international climate regime. The Paris Agreement of 2015 placed nationally determined mitigation and adaptation commitments within a shared temperature goal. More recent climate conferences have increasingly addressed how to finance implementation, respond to unavoidable damage, and make energy transitions equitable (United Nations, n.d.-a, n.d.-b; UNFCCC, n.d.-a).

Ethiopia's selection to host COP32 gives this evolution particular geographic and political significance. A decision adopted at COP30 accepts Ethiopia's offer to host the conference in Addis Ababa from 8 to 19 November 2027 and calls for a host-country agreement. The venue and dates are established in the decision; the substantive negotiating outcomes of COP32, which lies in the future, cannot yet be reported (UNFCCC, 2025).

Table 3.1. How international environmental priorities have evolved

Milestone	Core emphasis	Why it matters for Ethiopia
Stockholm, 1972	Pollution, environmental management, human well-being	Environmental health and resource stewardship
Rio Earth Summit, 1992	Sustainable development; climate and biodiversity conventions	Land, forests, water, development and global obligations
Kyoto, 1997	Binding emissions targets for specified industrialized countries	Differentiated responsibilities and carbon accounting
Paris, 2015	Universal climate framework; nationally determined contributions	Mitigation and adaptation planning within development priorities
COP27, 2022	Loss-and-damage funding arrangements	Recognition of impacts beyond feasible adaptation
COP28, 2023	First global stocktake; transition away from fossil fuels	Energy, resilience, ecosystem protection and implementation
COP29, 2024	New climate-finance goal of at least \$300 billion/year by 2035	Access, adequacy, terms and delivery of finance
COP30, 2025	Implementation, just transition, adaptation-finance emphasis	Connecting negotiated commitments to delivery
COP32, Addis Ababa, 2027	Scheduled future negotiating session; agenda/outcomes not yet determined	Opportunity to foreground African and Ethiopian priorities

Sources: United Nations (n.d.-a, n.d.-b); UNFCCC (2025, n.d.-a, n.d.-b). Conference themes are analytical summaries, not claims that any one topic was exclusive to a given year.

The terms fundamental, prominent, and urgent are useful analytical categories, not formal UN classifications. Fundamental concerns—safe water, clean air, healthy soils, biodiversity, and sanitation—remain necessary regardless of diplomatic cycles. Prominence describes the visibility of

an issue in treaties and negotiations; urgency describes the speed and severity of harm or the narrowing opportunity to prevent it. A drought can be locally urgent even when global negotiations focus on finance. Climate change increasingly amplifies older concerns without replacing them.

3.1 Concepts, pathways, and metrics

Population–environment relationships operate through multiple causal pathways. The number of people matters for aggregate demand, but environmental impacts depend just as strongly on consumption, production technology, land governance, settlement form, and ecological conditions. Population growth is therefore neither a sufficient explanation for degradation nor an excuse to overlook unequal resource use. The original chapter’s IPAT and Kaya identities remain valuable as accounting frameworks, provided they are not misinterpreted as causal models.

Accounting identities and interpretation

IPAT: $I = P \times A \times T$, where I denotes an environmental impact, P population, A affluence or output per person, and T impact per unit of output. The Kaya identity specializes this logic to energy-related carbon dioxide: $CO_2 = P \times (GDP/P) \times (Energy/GDP) \times (CO_2/Energy)$. Each term is measurable in principle, but observed relationships require consistent definitions, reference years, and sectoral boundaries.

A country may increase total energy demand as its population and economy expand while reducing emissions per unit of GDP through cleaner generation or efficiency. Conversely, low emissions per person do not imply low exposure to environmental harm. Household smoke, unsafe water, degraded soils, and flood risk can impose heavy health burdens in low-emitting communities. For Ethiopia, the correct analytical unit changes with the question: a watershed for water stress, a household for cooking fuels, a woreda for local service exposure, and national totals for emissions accounting.

Table 3.2. Priority indicators and their denominators

Domain	Preferred measure	Interpretive safeguard
Population pressure	Residents; households; population per inhabited km ²	Use harmonized boundaries; distinguish scenario from census
Land conversion	Hectares converted/year; share by original land class	Forest-cover loss is not automatically deforestation
Water security	Withdrawals / renewable resources by basin and season	National averages conceal basin-level scarcity
Energy transition	Clean-cooking access; electricity access; emissions intensity	Access, reliability and fuel stacking are distinct
Climate exposure	People within a validated hazard footprint	Exposure is not equivalent to vulnerability or losses
Environmental health	Population-weighted PM _{2.5} ; drinking-water safety	Use measured/modelled exposure with uncertainty

An important lesson from Chapter 2 is the need to distinguish geographical reconstruction from observation. The author’s 2025 regional reconstruction totals 127,935,000 across 14 current regions

plus ET99, a separately counted analytical geography. That benchmark is not a national census count and is not directly interchangeable with international population series. The World Bank’s World Development Indicators series, for example, reports 132,059,767 for 2024 and 135,472,051 for 2025. Differences in demographic definitions, models, timing, and geographic treatment must be reconciled before calculating rates (World Bank, 2026).




Figure 3.1. Ethiopia population estimates, 2023–2025 (World Bank World Development Indicators, as published in 2026). These are modeled international estimates, not census enumerations or the author’s regional reconstruction. Source: World Bank via FRED (2026).

3.2 Demographic change and environmental pressures

Demographic change influences the scale and geography of resource demand through population growth, age composition, urbanization, migration, and household formation. The same national population can produce very different environmental outcomes under different settlement and energy systems. Concentrated urban growth may reduce travel distances and infrastructure costs per resident, yet unmanaged expansion can increase flood exposure, air pollution, wastewater discharge, and pressure on peri-urban farmland. Rural population growth can intensify cultivation where land is scarce, but outcomes depend on tenure, agricultural productivity, and off-farm opportunities.

For Ethiopia, population growth is best treated as a denominator and planning driver, not a stand-alone measure of environmental degradation. A growing number of households may increase demand for dwellings, water connections, cooking devices, and waste services even where household size falls. At the same time, expanding access to electricity and clean cooking can reduce exposure to household air pollution. To quantify these pathways, analysts should pair observed household and population series with energy, land, water, and service indicators from matching years.

The author’s 2025 chapter included an illustrative table of population, urbanization, and mean household size from 1990 to 2035, alongside a synthetic “environmental pressure index.” Those values are not reproduced as empirical trends. The replacement figure above uses a traceable World Bank series and supports only the narrower claim of recent population growth. It does not measure ecological pressure or establish its causes.

Ethiopia and the global distribution of responsibility

Environmental responsibility and vulnerability are unevenly distributed. Carbon emissions are driven by energy systems, consumption, industrial production, and technology, while the costs of climate variability may fall disproportionately on rain-fed farming and pastoral communities. The distinction between total emissions, per-capita emissions, historical cumulative emissions, and exposure to climate hazards is essential to fair comparison. For COP32, these different metrics matter to discussions of mitigation ambition, adaptation needs, finance, and equity; none should be collapsed into a single ranking.

3.3 Land-use and land-cover change, forests, and restoration

Land cover describes what is physically present—forest, cropland, grassland, wetland, or built-up surfaces—while land use describes the human purpose to which land is put. A change in tree cover is not automatically equivalent to permanent deforestation: fire, harvest cycles, plantations, agroforestry, and forest degradation complicate interpretation. Satellite-based tree-cover-loss products are invaluable for identifying possible change, but land-cover transitions should be validated against classification rules, imagery, and field evidence before being attributed to agricultural expansion or other drivers.

In Ethiopia’s highlands, cultivated landscapes, steep slopes, and densely settled rural districts create particular concerns about soil erosion, watershed condition, and land fragmentation. In lowland regions, rangeland conversion, irrigated agriculture, woody-vegetation change, and settlement corridors pose different questions. A uniform national forest-loss narrative would obscure these differences. Restoration may improve infiltration, soil stability, habitat connectivity, and local livelihoods, but effectiveness depends on species selection, tenure, maintenance, and the ecological suitability of the intervention.

A credible empirical update would use an explicitly named and dated land-cover product, report hectares and percentages by region, apply a common classification crosswalk, and quantify uncertainty. It should distinguish gross loss from net change and tree-cover change from natural-forest conversion. Until those steps are complete, the original chapter’s illustrative forest-fragmentation and vegetation-anomaly time series should remain excluded.

A COP32 question: what constitutes verifiable restoration?

The Ethiopian contribution to international environmental discussion can be strengthened by moving from counts of seedlings or hectares reportedly treated toward durable outcomes: survival rates, changes in soil cover, erosion, water availability, biodiversity, and benefits to communities. Remote sensing, repeat field sampling, and locally grounded monitoring are complementary rather than interchangeable. Carbon accounting should not displace the equally important objectives of livelihoods, ecosystem health, and resilience.

3.4 Agriculture, pastoral systems, and rangeland dynamics

Agriculture and pastoralism connect demographic change directly to climate, water, soils, and food systems. Highland mixed farming and lowland livestock systems differ in their use of land and mobility, and neither can be adequately assessed through a single stocking-density or vegetation index. In pastoral areas, seasonal movement can be an adaptive strategy that matches livestock to spatially variable forage and water. Restricting mobility may increase localized pressure even when aggregate herd numbers are unchanged.

Afar and Somali illustrate why regional averages require careful interpretation. Using the author's 2025 reconstructed populations and accompanying land areas, their arithmetic densities are approximately 26 and 28 people per km², respectively. These figures describe residents per unit of administrative area; they do not estimate the number of pastoralists, seasonal occupancy, herd size, forage supply, or the sustainable stocking rate. Nor do they imply that people and livestock are evenly distributed across the territory.

Rainfall anomalies, normalized difference vegetation index (NDVI), livestock inventories, water-point functionality, market prices, and participatory mobility information can provide a fuller picture when observed for matching seasons and geographies. NDVI is a proxy for greenness, not a direct measure of edible forage. Market offtake may reflect normal commercialization or distress sales; interpretation requires prices, herd composition, and context. In conflict-affected settings, precise mapping of routes and watering points may also create protection risks.

From environmental stress to adaptation capacity

Policy options include protecting mobility corridors, strengthening veterinary services, investing in reliable water access, improving market connectivity, and providing early warning and shock-responsive support. Their effectiveness depends on local institutions and land rights. The practical COP32 lesson is that adaptation is not merely a technological intervention: it can involve preserving the flexibility of existing livelihoods and ensuring that climate finance reaches communities exposed to recurrent hazards.

Part I synthesis and evidence audit

The central proposition is that environmental outcomes emerge from interactions among people, consumption, technology, institutions, and ecosystems. Population growth changes the scale of demand; land governance and livelihood systems shape how demand translates into environmental pressure; climate hazards affect people unequally. The path from Stockholm to Addis Ababa shows a widening policy agenda, with implementation and distributional fairness increasingly prominent alongside enduring conservation and public-health concerns.

Evidence status	Treatment in this revision
Verified institutional history	Stockholm, Rio, Paris and major COP milestones summarized from UN/UNFCCC sources
Confirmed future event	COP32 Addis Ababa, 8–19 November 2027; outcomes not anticipated
Observed/modelled external series	World Bank population estimates for 2023–2025; explicitly sourced
Author reconstruction	127,935,000 (2025), including ET99; scenario only, not official count
Excluded placeholders	Synthetic pressure, forest, NDVI, mobility, and rangeland graphs from 2025 PDF

PART II. 3.5 Water resources, irrigation, and watersheds

Ethiopia’s water geography is structured by pronounced contrasts among river basins, seasons, and livelihoods. The Abbay (Blue Nile), Awash, Omo–Gibe, and Baro–Akobo systems differ in hydrology, population settlement, irrigation potential, and downstream connections. National totals for renewable water resources cannot reveal scarcity at a particular intake, aquifer, or dry-season reach. Water stress must therefore be assessed at basin or sub-basin scale, with withdrawals and environmentally necessary flows referenced to comparable periods.

The demographic pathway is not simply “more people, less water.” Population growth increases aggregate demand for drinking water, sanitation, food, and energy, but per-capita demand also responds to urbanization, irrigation technology, industrialization, and pricing. Irrigation can improve food-system resilience while increasing competition for water where withdrawals are poorly measured. Likewise, reservoir capacity does not by itself establish year-round service reliability, ecological sustainability, or equitable allocation.

In the highlands, watershed degradation and erosion can affect sediment delivery, reservoir operations, and agricultural productivity. In the lowlands, episodic floods and droughts may dominate seasonal availability. Groundwater can buffer variability but requires monitoring of recharge, extraction, salinity, and water quality. Remote sensing, river gauges, household surveys, and irrigation registries offer complementary—not interchangeable—measurements.

Analytical priorities for Ethiopia

A decision-ready basin assessment should report renewable supply and actual withdrawals by season; irrigated area and crop-water use; groundwater trends; environmental-flow compliance; and population with reliable, safely managed water. For cross-border basins, hydrological evidence and cooperative institutions matter together. Claims about national water abundance should not be extrapolated to local water security.

3.6 Climate variability and change: exposure, sensitivity, and adaptation

Climate risk emerges from the interaction of a hazard with exposed people and assets, their sensitivity, and their capacity to respond. A flood map is not a map of losses; a rainfall anomaly is not a direct measure of hunger; and a warming trend does not determine the impact on every community. In Ethiopia, drought-prone pastoral lowlands, rain-fed farming highlands, rapidly growing cities, and flood-exposed river corridors face distinct combinations of risk.

A warming climate can intensify heat exposure and alter water demand, while changing rainfall variability complicates planting decisions, pasture availability, and reservoir management. The strength, direction, and timing of precipitation changes vary by region and season; therefore, an Ethiopia-wide single rainfall trajectory is insufficient. Extreme-event attribution, long-term trend estimation, and near-term forecasting answer different questions and should not be conflated.

For public health, climate-sensitive pathways include heat illness, waterborne disease after flooding, disrupted maternal and child health services, changes in vector ecology, and food insecurity. The relevant denominator is often the number of people living or working within a hazard footprint, disaggregated by age, disability, livelihood, and service access—not merely the land area affected.

From adaptation commitments to implementation

As COP32 approaches, adaptation should be evaluated through operational outcomes: timely early-warning dissemination, maintained all-weather access to health facilities, flood-resilient water systems, drought preparedness, and social protection that can scale before households exhaust assets. Financing announcements are inputs; functioning systems and reduced vulnerability are outcomes. A rigorous Ethiopian COP32 narrative should distinguish these levels.

3.7 Food security, nutrition, and environmental shocks

Food insecurity is a multidimensional outcome of production, prices, income, access, and stability. Rainfall deficits can reduce yields or forage, but conflict, transport interruptions, market prices, and household purchasing power can magnify or offset their effects. Ethiopia's mixed farming and pastoral systems consequently require different surveillance and response designs. A vegetation index may flag forage stress without directly measuring livestock condition, household consumption, or child malnutrition.

The Integrated Food Security Phase Classification (IPC) describes acute food insecurity using convergent evidence. Its categories are not interchangeable with childhood stunting or wasting, which are measured on different populations, timescales, and biological criteria. Likewise, “people in need,” “people targeted,” and “people reached” are distinct humanitarian quantities. Reports should preserve these distinctions and their observation dates.

Nutrition outcomes arise through both direct and indirect pathways: reduced dietary diversity, infectious disease, compromised water and sanitation, lost household income, and disrupted care. Young children, pregnant and lactating women, displaced households, and people with limited market access may face disproportionate risk. A geographically precise account would connect

nutrition survey outcomes to livelihood zones, food prices, access constraints, and documented shocks while avoiding ecological inference from region-level averages.

COP32 implication: resilience as a measurable food-system outcome

An actionable agenda would link forecast-based financing, irrigation and water harvesting, livestock mobility safeguards, market access, nutrition services, and social protection. The policy question is not merely whether climate adaptation is funded, but whether investments reduce the probability that an environmental shock becomes a nutritional crisis.

3.8 Urbanization, air quality, and environmental health

Urban environmental health combines outdoor pollution from transport, industry, dust, and waste burning with household exposure from cooking and heating fuels. PM_{2.5} denotes fine particulate matter; NO₂ is nitrogen dioxide. Their concentrations are not interchangeable, and modeled city averages should not be interpreted as individual exposure. Household air pollution can be severe even in communities with relatively little motor traffic.

WHO reports that about 2.1 billion people worldwide still relied on polluting fuels and technologies for cooking, and estimates 2.9 million premature deaths attributable to household air pollution in 2021. These are global estimates, not Ethiopia-specific counts (WHO, 2025). For Ethiopia, the WHO country profile reports 7% of the population primarily relying on clean cooking fuels and technologies in 2022 (uncertainty interval 3%–13%). The year and uncertainty interval are essential parts of that estimate.

These indicators are relevant to equity: rural households may face limited clean-fuel availability, while low-income urban and peri-urban residents can face combined household and ambient exposures. A 2016 Ethiopia DHS secondary analysis published in the Ethiopian Journal of Health Development estimated cooking-smoke exposure among 61.4% of its weighted analytical household sample; this is a study-specific exposure definition and must not be substituted for WHO’s clean-fuel-use indicator.

A verified empirical snapshot

Indicator	Observed/modelled estimate	Scope and reference year
Primary reliance on clean cooking fuels	7% (uncertainty interval 3%–13%)	Ethiopia; WHO country profile, 2022
Urban annual mean PM_{2.5}	20.87 µg/m ³ (14.19–31.75)	Ethiopia; WHO country profile, 2019
Cooking-smoke exposure	61.4% (95% CI 59.6–63.2)	Ethiopia; weighted 2016 DHS study sample
Household air pollution mortality	2.9 million deaths	Global WHO estimate, 2021

Note. Indicators have different populations, definitions, observation years, and estimation methods; they must not be plotted as a single trend or treated as directly comparable rates.

The WHO country profile also reports an age-standardized mortality rate attributable to ambient and household air pollution of 120.6 per 100,000 in Ethiopia in 2019 (104.3–137.3). This modeled combined-exposure burden is not a count of deaths attributable to household cooking smoke alone. Cleaner cooking, emissions standards, urban transport planning, waste management, and reliable monitoring all have potential health and climate co-benefits, but benefits must be evaluated rather than assumed.

PART II synthesis: an Ethiopian environmental-health agenda for COP32

The four domains converge around a common planning principle: environmental stress becomes a demographic and health crisis through unequal exposure, livelihood dependence, inadequate infrastructure, and limited adaptive capacity. Water security, resilient food systems, climate-ready health services, and cleaner household energy are therefore not peripheral “co-benefits” of climate policy. They are central measures of whether climate action improves human wellbeing.

The historical shift from environmental protection toward adaptation, equity, and implementation is particularly consequential for Ethiopia. Hosting COP32 creates an opportunity to present measurable national priorities, request appropriately targeted finance, and contribute African evidence on climate vulnerability and locally effective responses. No outcome of the 2027 conference is presumed in this October 2026 revision.

PART III · Energy, Biodiversity, Disaster Risk, and Infrastructure

3.9 Energy Transitions and Household Fuels

Energy transitions have two distinct dimensions in Ethiopia: the composition of electricity generation and the fuels actually used by households and firms. A predominantly renewable power-generation fleet does not mean that households enjoy reliable electricity or clean cooking. This distinction is central to both environmental accounting and health policy. Household reliance on wood, charcoal, dung, and crop residues links energy poverty to indoor particulate exposure, time spent collecting fuel, pressure on woody biomass, and gendered labor. The size of each pathway must be established empirically rather than inferred from a national electricity-generation statistic.

Ethiopia’s National Energy Compact reports that in 2025 installed generation capacity was 9,761 MW, more than 95% from renewable sources, mainly hydropower. The same compact distinguishes a broad measure of electricity access (65% in 2025, including some off-grid solutions) from basic Tier 1-or-higher service (44%). These measures have different definitions and should not be plotted as a single comparable series. The compact also reports continued extensive use of firewood for cooking in 2024. The energy-access problem is therefore not solved by adding generation capacity alone: affordability, distribution, service quality, and appliance availability matter (World Bank, 2026).

Hydropower creates an opportunity for low-carbon electrification, but it also exposes electricity supply to hydrological variability and entails basin-scale trade-offs. Solar, wind, geothermal, storage,

and resilient transmission can diversify the supply portfolio. Decentralized systems may be appropriate for dispersed settlements, but must be evaluated for lifecycle cost, maintenance, and actual delivered service. A just transition should track whether low-income and rural households gain reliable power and cleaner cooking—not merely the number of megawatts installed.

A COP32 question: whose transition, and measured how?

At COP32, Ethiopia can make the case that mitigation and development are not competing objectives when financing supports dependable clean electricity, modern cooking, and resilient grids. Credible reporting should separate installed capacity (MW), electricity generated (GWh), household connection, minimum service tier, clean-cooking use, and affordability. It should also report distributional differences across rural and urban populations, rather than treating national averages as universal access.

3.10 Biodiversity, Protected Areas, and Human Pressure

Ethiopia’s ecological diversity extends from Afroalpine environments and highland forests to drylands, wetlands, and Rift Valley ecosystems. Biodiversity loss arises from multiple interacting pressures: land conversion, habitat fragmentation, unsustainable resource extraction, invasive species, pollution, and climate change. Population growth can amplify land and resource demand, but it is not a sufficient causal explanation. Tenure arrangements, livelihood alternatives, infrastructure placement, enforcement, and consumption patterns determine how demographic change translates into ecological pressure.

The global biodiversity assessment published by IPBES (2019) estimated that approximately one million animal and plant species face extinction risk worldwide. That figure is global, not an estimate for Ethiopia. It nevertheless clarifies why biodiversity has moved from a conservation concern to a foundational issue for food systems, water regulation, disease ecology, and economic resilience. In Ethiopia, the relevant empirical questions concern land-cover transitions, habitat connectivity, species status, and the distribution of benefits and costs from conservation.

Protected-area boundaries are not a substitute for ecological effectiveness. A meaningful evaluation would overlay verified protected-area polygons with consistent land-cover classifications, vegetation condition, infrastructure corridors, and locally validated ecological surveys. For communities living adjacent to protected landscapes, conservation design should account for grazing, water access, compensation, and conflict prevention. The appropriate policy objective is durable ecosystem protection alongside legitimate livelihood needs.

From separate conventions to integrated stewardship

The 1992 Rio Earth Summit institutionalized a wider agenda linking climate, biological diversity, and sustainable development. In contemporary policy, these agendas increasingly intersect: forests store carbon and sustain habitat; wetlands attenuate flooding; intact rangelands support livelihoods and ecological connectivity. Yet a carbon-focused restoration program is not automatically biodiversity-positive. Species composition, water demand, ecological suitability, and community participation must be assessed before claiming co-benefits.

3.11 Disasters, Hazards, and Ecosystem Services

Environmental hazards become disasters through the interaction of physical events with exposed people, vulnerable assets, and limited capacity to anticipate and respond. Floods, droughts, landslides, wildfire, and pest outbreaks have different causal mechanisms, yet can produce cascading effects across transport, agriculture, food prices, drinking water, and health services. Risk therefore cannot be summarized by the frequency of a hazard alone. A high-intensity flood in an uninhabited catchment has different consequences from a smaller flood affecting a densely settled, poorly drained neighborhood.

The United Nations Office for Disaster Risk Reduction (UNDRR, 2025) estimates that global annual disaster costs exceed US\$2.3 trillion when indirect and ecosystem impacts are included. This is a modeled global burden, not an Ethiopian loss estimate, and is not directly comparable with routinely reported direct damages. The key policy lesson is that investments in prevention, risk-sensitive land use, early warning, and resilient services can avert losses that are missed by narrow post-disaster accounting.

Ecosystems can moderate—but cannot eliminate—hazard impacts. Restored upper catchments may reduce erosion and sediment delivery; wetlands and floodplains can store water; urban trees and permeable surfaces may moderate heat and runoff. Their effects are place-specific, conditional on rainfall, soil, slope, land management, and maintenance. Nature-based measures should be tested against engineering alternatives and combined with drainage, emergency response, health-system preparedness, and social protection where appropriate.

Population exposure is not the same as disaster loss

An empirical Ethiopian disaster-risk atlas would distinguish (1) a validated hazard footprint, (2) the population and assets located within it, (3) social and infrastructural vulnerability, and (4) observed losses. Population grids must be dated and harmonized to the hazard period; displacement and seasonal pastoral movement require separate treatment. Climate-attribution statements need additional analysis and cannot be inferred from a single extreme event.

3.12 Infrastructure, Mining, and Environmental Footprints

Roads, dams, industrial parks, transmission lines, urban expansion, and mineral extraction can generate substantial economic benefits while changing landscapes, hydrology, and settlement patterns. Infrastructure may open markets and expand services, but also fragment habitats, alter sediment transport, increase construction emissions, and expose neighboring communities to air, water, or noise pollution. Environmental assessment should therefore address the full project life cycle: siting, construction, operation, decommissioning, and cumulative effects across connected projects.

Environmental impact assessment (EIA) is most valuable before a project location and design become irreversible. Alternatives analysis should compare not only engineering cost but also resettlement, ecological sensitivity, downstream effects, and social distribution of risks. Strategic environmental assessment is especially important when a series of roads, energy installations, or

extraction sites transforms a corridor or basin. A project-by-project review may miss the cumulative footprint of the entire development program.

For hydropower, the relevant questions extend beyond low operational carbon intensity: environmental flows, sediment, fisheries, upstream and downstream livelihoods, reservoir impacts, and transboundary cooperation all matter. For mining, priority safeguards include tailings stability, water quality, waste handling, rehabilitation bonds, occupational health, and transparent community consultation. For urban and industrial infrastructure, air emissions, wastewater treatment, and climate resilience should be assessed jointly.

COP32: from project approval to measurable safeguards

Hosting COP32 provides an opportunity to link climate-finance claims with project-level accountability. Financing should specify measurable outcomes, safeguards, baseline conditions, independent monitoring, grievance mechanisms, and benefits for affected populations. Infrastructure that is low-carbon but environmentally or socially damaging should not be described as unconditionally sustainable.

PART III Synthesis: Four Connected Transitions

The four sections describe linked transitions in energy, ecosystems, risk governance, and physical infrastructure. Their common challenge is measurement. Installed renewable capacity is not clean-cooking access; protected-area designation is not ecological integrity; hazard occurrence is not disaster loss; and project approval is not environmental performance. For Ethiopia's COP32 agenda, the strongest contribution will be to translate these distinctions into transparent investment criteria and outcome indicators, connecting global climate commitments to locally demonstrable public benefits.

Table 3.3. Evidence and measurement priorities for PART III

Domain	Defensible indicator	Key interpretive caution
Energy	Electricity service tier; clean cooking; generation by source	Generation share $\neq$ household energy access
Biodiversity	Habitat change; connectivity; verified species status	Global IPBES estimate $\neq$ Ethiopia count
Disaster risk	Population exposed; losses; resilience investment	Hazard frequency $\neq$ realized loss
Infrastructure	EIA compliance; water/air monitoring; restoration	Project approval $\neq$ sustained performance

Note. This table is an analytical measurement framework, not a dataset of observed Ethiopian outcomes. No synthetic charts from the 2025 manuscript are reproduced.

PART IV — Environmental governance and the road to COP32

3.13 Waste, Sanitation, and Circularity

Population growth, urban expansion, and changing consumption patterns increase the complexity of municipal waste management. The principal environmental concern is not merely the volume of waste generated, but whether waste is prevented, collected, separated, treated, recycled, or disposed of safely. Uncollected solid waste can obstruct drainage and aggravate flood exposure; open burning can contribute to air pollution; unmanaged organic waste can produce methane; and inadequate fecal-sludge management can contaminate water sources. These pathways link environmental management directly to public health and urban resilience.

A circular-economy approach moves beyond end-of-pipe disposal toward waste avoidance, material reuse, repair, recycling, and recovery, with safeguards for workers and communities. Globally, UNEP (2024) estimated municipal solid waste generation at 2.1 billion tonnes in 2023 and projected 3.8 billion tonnes by 2050 under its modeled trajectory. The latter is a projection, not an observed quantity. Neither figure should be treated as an Ethiopian estimate.

Ethiopia's NDC 3.0 identifies urban waste services and climate-resilient infrastructure as implementation priorities. For Ethiopian cities, a credible monitoring system should separately report collection coverage, safe treatment or disposal, recycling, waste composition, landfill methane management, safely managed sanitation, and the spatial distribution of underserved neighborhoods. Administrative coverage claims require clear definitions and independent quality checks.

Environmental-health pathways and measurement

Pathway	Priority indicator	Interpretive safeguard
Uncollected waste and flooding	Collection coverage; drainage blockage reports	Collection is not equivalent to safe disposal
Organic waste and methane	Waste composition; controlled treatment; methane capture	Do not infer emissions reductions from collection alone
Unsafe sanitation and water contamination	Safely managed sanitation; fecal-sludge treatment	Distinguish improved facilities from safely managed services
Informal waste labor	Occupational injury, protective equipment, income	Account for informal workers and gendered risks

3.14 Gender, Equity, and Environmental Justice

Environmental hazards are not experienced uniformly. Exposure differs by place, occupation, age, disability, housing quality, wealth, and access to infrastructure. Women and girls may bear disproportionate time burdens for collecting water and cooking fuels, while children, older adults, outdoor workers, and people with chronic illnesses may be especially vulnerable to heat and air pollution. Such patterns require empirical verification rather than universal assumptions about every Ethiopian community.

Environmental justice adds a distributional and procedural dimension to environmental policy. Distributional justice asks who bears pollution and climate risks and who receives investments. Procedural justice asks who participates in decisions about land, water, energy, resettlement, and climate adaptation. Recognition justice asks whether pastoralist, displaced, low-income, and other marginalized populations are visible in the data and decision-making processes. Climate interventions can improve aggregate outcomes yet deepen inequities if land tenure, livelihoods, or access to services are overlooked.

For Ethiopia, equity-sensitive analysis should combine disaggregated household surveys with geographically explicit hazard and service-access measures. It should report exposure and vulnerability separately: living in a drought-prone area is not the same as experiencing a drought-related loss. National averages should not obscure regional or within-city disparities, and analyses should avoid publishing identifiable locations of vulnerable households.

Practical equity tests for proposed interventions

Every major project should ask: Who is exposed before intervention? Who receives benefits? Who pays costs or loses access? Who participates in decisions? How will unintended harms be monitored? This framework is particularly relevant to watershed restoration, urban waste infrastructure, renewable-energy projects, and pastoral rangeland management.

3.15 Governance, Policy, and Financing for Sustainability

Ethiopia's environmental governance has evolved from sector-specific natural-resource management toward integrated climate planning. Key frameworks include the Climate-Resilient Green Economy strategy, the National Adaptation Plan, successive nationally determined contributions (NDCs), and the Long-Term Low-Emissions Development Strategy. These instruments have different functions: strategies articulate direction, NDCs communicate commitments under the Paris Agreement, adaptation plans prioritize resilience measures, and monitoring systems assess implementation.

Ethiopia submitted NDC 3.0 to the UNFCCC on 26 September 2025. It specifies a 70.3% greenhouse-gas reduction by 2035 against a revised business-as-usual reference scenario. This is a scenario-relative target—not a 70.3% decline from a measured historical emissions level—and cannot be interpreted as an achieved reduction. The NDC distinguishes actions funded through domestic effort from conditional actions dependent on external support (Federal Democratic Republic of Ethiopia, 2025).

The NDC estimates financing needs of approximately US\$106.35 billion for 2025–2035, including US\$66.35 billion for mitigation and US\$40 billion for adaptation. These are planned investment requirements, not funding already received. A useful COP32-facing assessment would distinguish pledged finance, committed finance, disbursed finance, and independently verified expenditure, alongside the terms of financing and local distribution of benefits.

Governance question	Evidence needed	Why it matters
Are targets being implemented?	Annual, sector-specific indicators and baselines	Targets and outcomes are not interchangeable

Is finance reaching projects?	Commitments, disbursements, grants/loans, execution	Announcements do not establish delivery
Who benefits?	Regional and socioeconomic disaggregation	National totals can hide exclusion
Are reductions additional and durable?	Transparent inventories, verification, permanence tests	Avoid double counting and overclaiming
Can institutions coordinate?	Clear mandates and interoperable reporting	Fragmentation impedes delivery

COP32: Ethiopia as host and negotiating participant

The UNFCCC confirms that parties accepted Ethiopia’s offer to host COP32 in 2027. The conference is scheduled in Addis Ababa for 8–19 November 2027. Hosting provides visibility and convening opportunities, but does not by itself guarantee financing, favorable negotiating outcomes, or domestic implementation. The analytically important question is whether preparations produce durable systems for adaptation planning, climate finance tracking, transparent emissions inventories, and locally accountable delivery.

COP32 also offers a useful historical lens: international environmental diplomacy has progressed from pollution control and conservation to sustainable development, emissions mitigation, adaptation, loss and damage, and the practical challenge of implementation. These are cumulative priorities rather than successive replacements. Ethiopia’s experience with drought, land restoration, urbanization, and energy access illustrates why they must be addressed together.

3.16 Data, Methods, and Indicators

A population–environment monitoring system should integrate demographic denominators, earth observation, environmental monitoring, administrative records, and household surveys. Population counts must be aligned by reference year and geographic boundaries. Climate observations must be distinguished from modeled reanalyses and future scenarios; satellite proxies from ground-validated measurements; and administrative service records from population coverage estimates.

Recommended inputs include Ethiopian statistical and sectoral data; EDHS and other household surveys; CHIRPS precipitation; ERA5 climate reanalysis; satellite land-cover and vegetation products; water-basin measurements; WHO environmental-health indicators; and UNFCCC greenhouse-gas reporting. Each indicator should carry source, geographic scale, time period, definition, denominator, uncertainty, and revision history.

Proposed Ethiopia population–environment indicator registry

Domain	Illustrative indicator	Essential metadata
Climate exposure	Population exposed to heat, drought, or flood hazard	Hazard threshold, grid year, population vintage

Land and forests	Land-cover transitions; restoration survival	Classification, sensor, validation accuracy
Water and sanitation	Safe water access; safely managed sanitation	Survey definition, season, residence
Air quality	Population-weighted PM _{2.5} exposure	Monitor/model source, calibration, uncertainty
Energy	Clean-cooking access; electricity reliability	Fuel-use definition, service continuity
Finance	Adaptation funding disbursed per vulnerable resident	Funding stage, currency year, denominator

Methodological rule: no fabricated trend lines, synthetic observations, or unverified regional rankings. Where evidence is incomplete, use a conceptual diagram, an indicator specification, or an explicit data gap rather than a numerical graphic. Geographic analysis should document harmonization across Ethiopia's changing administrative boundaries and avoid treating modeled small-area estimates as census enumerations.

3.17 Case Studies and Ethiopian Successes

Ethiopia offers important examples of environmental initiatives, but a defensible case study distinguishes implementation scale from measured ecological or health outcomes. The Green Legacy Initiative, highland watershed restoration, community-based natural-resource management, clean-energy expansion, and pastoral water-management programs are candidates for evaluation—not automatic evidence of success on every outcome.

Case study A — Landscape restoration and the Green Legacy Initiative

Tree-planting campaigns and restoration initiatives may contribute to erosion control, habitat recovery, local cooling, and carbon sequestration. However, the number of seedlings distributed or planted is an activity measure, not a verified estimate of surviving forest, carbon storage, or biodiversity improvement. Evaluation requires survival rates, species suitability, land-tenure safeguards, baseline land cover, and multi-year satellite and field measurements. Claims about forest-cover change should be checked for differences in forest definitions and observation methods.

Case study B — Climate-resilient watersheds and rural livelihoods

Highland watershed interventions can combine soil conservation, terracing, vegetation recovery, and water management. Plausible benefits include reduced erosion, improved infiltration, and more reliable agricultural production, but their magnitude is location-specific. Rigorous evaluation compares treated and comparable untreated catchments over sufficient periods while accounting for rainfall variation, land use, and upstream–downstream effects.

Case study C — Urban clean energy and environmental health

Renewable electricity generation is an important national asset, yet electricity supply alone does not establish household adoption of clean cooking or reductions in indoor smoke exposure. Case studies should measure household fuel stacking, affordability, reliability, ventilation, and health-relevant

exposure. Distributional analysis is essential to determine whether benefits reach poorer households and secondary cities.

Case study D — Pastoral adaptation and mobility

In lowland pastoral systems, seasonal movement can be a rational adaptation to variable forage and water. Interventions should assess access to mobility corridors, conflict risks, animal health, and market functioning rather than equating sedentarization with environmental improvement. Participatory evaluation and protection of sensitive location data are essential.

PART IV synthesis — From hosting to implementation

Ethiopia’s COP32 moment is most consequential when viewed not as a single diplomatic event but as a test of institutional capacity: can environmental priorities be translated into financed, geographically targeted, socially equitable, and independently measured outcomes? The central lesson of Chapter 3 is that population and environment are linked through land, water, energy, livelihoods, health, and governance. Sustainable progress requires attention to each linkage, with special protection for those least able to absorb environmental shocks.

The global environmental agenda has broadened over five decades without abandoning its original foundations. Clean air, safe water, healthy ecosystems, and human well-being remain fundamental; climate mitigation, adaptation, and finance have become prominent; and implementation, equity, and loss-and-damage responses have become urgent. Addis Ababa in 2027 provides a compelling setting in which these strands can converge.

Chapter 3 | References

- Atnafu, D. D., Ketemaw, A., Menber, Y., Wassihun, Y., & Guadie, H. A. (2021). *Cooking smoke (household air pollution) exposure status of households in Ethiopia: A further analysis based on 2016 DHS data*. *Ethiopian Journal of Health Development*, 35(3), 183–193. <https://ejol.aau.edu.et/index.php/EJHD/article/view/3815>
- Center for Climate and Energy Solutions. (2026). *Key negotiations and related outcomes of the UN climate change conference in Belém*. <https://www.c2es.org/wp-content/uploads/2026/02/COP30-Summary-FINAL5.pdf>
- Federal Democratic Republic of Ethiopia. (2025). *Ethiopia NDC 3.0*. United Nations Framework Convention on Climate Change. <https://unfccc.int/sites/default/files/2025-09/Ethiopia%20NDC%203.0%20Final.pdf>
- Food and Agriculture Organization of the United Nations. (n.d.). *AQUASTAT: FAO’s global information system on water resources and agriculture*. <https://www.fao.org/aquastat/>
- Integrated Food Security Phase Classification. (n.d.). *IPC: Integrated Food Security Phase Classification*. <https://www.ipcinfo.org/>

- Intergovernmental Panel on Climate Change. (2022). *Climate change 2022: Impacts, adaptation and vulnerability*. Cambridge University Press. <https://www.ipcc.ch/report/ar6/wg2/>
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. (2019). *Global assessment report on biodiversity and ecosystem services* (E. S. Brondízio, J. Settele, S. Díaz, & H. T. Ngo, Eds.). <https://doi.org/10.5281/zenodo.3831673>
- International Energy Agency. (2026). *National Electrification Program 2.0 (NEP 2.0): Ethiopia*. <https://www.iea.org/policies/30385-national-electrification-program-20-nep-20>
- United Nations. (n.d.-a). *United Nations Conference on Environment and Development, Rio de Janeiro, Brazil, 3–14 June 1992*. <https://www.un.org/conferences/environment/rio1992>
- United Nations. (n.d.-b). *United Nations Conference on the Human Environment, Stockholm 1972*. <https://www.un.org/en/conferences/environment/stockholm1972>
- United Nations Economic Commission for Africa. (2026, September 9). *Stakeholders close CCDA-14 with strong calls for climate finance, stronger agency and delivery ahead of COP31*. <https://uneca.org/stories/stakeholders-close-ccda-14-with-strong-calls-for-climate-finance%2C-stronger-agency-and>
- United Nations Environment Programme. (2024). *Global waste management outlook 2024: Beyond an age of waste—Turning rubbish into a resource*. <https://www.unep.org/en/publications/report/global-waste-management-outlook-2024>
- United Nations Framework Convention on Climate Change. (2025a). *Dates and venues of future sessions (COP30 decision; advance unedited version)*. https://unfccc.int/sites/default/files/resource/COP30_2g_dates_venues_auv.pdf
- United Nations Framework Convention on Climate Change. (2025b). *UN Climate Change quarterly update: Q4 2025*. <https://www.unfccc.int/about-us/reports-highlights/quarterly-updates/un-climate-change-quarterly-update-q4-2025>
- United Nations Framework Convention on Climate Change. (n.d.-a). *Conference of the Parties (COP)*. <https://www.unfccc.int/process/bodies/supreme-bodies/conference-of-the-parties-cop>
- United Nations Framework Convention on Climate Change. (n.d.-b). *UN Climate Change: Who we are and what we do*. <https://unfccc.int/resource/bigpicture/index.html>
- United Nations Office for Disaster Risk Reduction. (2025). *Global assessment report 2025: Resilience pays—Financing and investing for our future*. <https://www.undrr.org/gar/gar2025>
- World Bank. (2026a). *Ethiopia national energy compact: Mission 300*. <https://www.worldbank.org/content/dam/theworldbankgroupasset/document/2026/Ethiopia-National-Energy-Compact-Mission-300.pdf>
- World Bank. (2026b). *Population, total for Ethiopia* [World Development Indicators; series POPTOTETA647NWDB]. Federal Reserve Bank of St. Louis. <https://fred.stlouisfed.org/series/POPTOTETA647NWDB>

- World Health Organization. (2018). *Opportunities for transition to clean household energy in Ethiopia: Application of the WHO Household Energy Assessment Rapid Tool (HEART)*.
<https://www.who.int/publications/i/item/9789241514491>
- World Health Organization. (2025, December 16). *Household air pollution*.
<https://www.who.int/news-room/fact-sheets/detail/household-air-pollution-and-health>
- World Health Organization. (n.d.-a). *Ethiopia: Country health data*.
<https://data.who.int/countries/231>
- World Health Organization. (n.d.-b). *Household air pollution: Global Health Observatory*.
<https://www.who.int/data/gho/data/themes/topics/topic-details/GHO/household-air-pollution>
- World Resources Institute. (2026). *Ethiopia's NDC 3.0: Climate ambition and development*.
<https://www.wri.org/technical-perspectives/ethiopia-ndc-3-climate-development>